



PRODUCT DATA SHEET

Ready-to-Screen Tissue BLOTS[™] Brain Tissue Region- Specific Blots - Single Species

Catalog #:	TB59		
Lot #:			
Components:	Protein blot of Primate Brain Region Tissue samples arranged as follows:		
	Lane 1 Protein Marker*	211,806	Myosin
	Lane 2 Cerebellar Cortex (Cerebellum) Lane 3 Hippocampus	121,020	β-galactosidase
	Lane 4 Striatum	100,216	Bovine Serum Albumin
	Lane 5 Thalamus	54,395	Ovalbumin
	Lane 6 Pons Lane 7 Medulla	34,393	Ovarbumin
	Lane 8 Hypothalamus	38,708 29,806	Carbonic Anhydrase Soybean Trypsin Inhibitor
	Lane 9 Temporal Lobe Lane 10 Amygdala	20,040	Lysozyme
	Lane 11 Occipital Lobe	7,331	Aprotinin
	Lane 12 Hippocampal Gyrus		
	Lane 13 Frontal Lobe Lane 14 Cingulate Gyrus	* Lot #: 300002325-BR	
	Lane 15 Parietal Lobe		

Size: 1 Blot

Storage Condition: 4° C

Methods Involved: The proteins were isolated from various primate brain region tissues by preparing a tissue homogenate in the presence of protease inhibitors. Protein samples (50μg) from each tissue were solubilized in SDS-lysis buffer and electrophoresed in a 15 well, 4-20% SDS-polyacrylamide gradient gel, followed by electroblotting on PVDF membrane.

<u>Quality Control</u>: Proteins isolated from each lot were run on 4-20% gel and stained with G-Biosciences *RapidStain*[™] to check for its quality. Actin antibody was used to test the separation and transfer of protein from each lot.

Instructions for Use: Remove the blot (membrane) from the pouch and wash with an appropriate buffer (1X TBST or PBST) 1-2 times. Block the membrane with a protein blocking agent; e.g., G-Biosciences NAP^{TM} -Blocker or BLOT- $QuickBlocker^{\text{TM}}$, and incubate with the primary and secondary antibodies diluted in blocking solution, following the standard protocol. Develop the blot with chemiluminescent or chromogenic detection reagents for the detection of the specific protein.

Rev 11.18.08-SA/MM/IA

