



290PR-02

*G-Biosciences* ♦ 1-800-628-7730 ♦ 1-314-991-6034 ♦ [technical@GBiosciences.com](mailto:technical@GBiosciences.com)

A Geno Technology, Inc. (USA) brand name

# EndotoxinOUT™ Resin

(Cat. # 786-367, 786-368, 786-369)



think proteins! think G-Biosciences [www.GBiosciences.com](http://www.GBiosciences.com)

## INTRODUCTION

EndotoxinOUT™ consists of 6% cross-linked agarose covalently linked to polymyxin B to bind and remove harmful pyrogens from a solution. Polymyxin B is a family, polymyxin B1 and B2, of antibiotics that bind to the negatively charged site of the lipid A portion of bacterial lipopolysaccharide layer neutralizing the endotoxic activity.

The covalent coupled agarose and polymyxin B is a stable matrix that resists leaching. An ideal product for the clean up of buffers, cell culture media, protein solutions, nucleic acid (DNA) samples and pharmacological components.

## ITEM(S) SUPPLIED

Cat. #	Description	Size *
786-367	EndotoxinOUT™	10ml resin
786-368	EndotoxinOUT™	1L resin
786-369	EndotoxinOUT™	5 x 1ml columns

\*EndotoxinOUT™ resin is supplied as a 50% slurry with 20% ethanol as a preservative.

## STORAGE CONDITIONS

It is shipped at ambient temperature. Upon arrival, store refrigerated at 4°C, **DO NOT FREEZE**. This product is stable for 1 year at 4°C.

## SPECIFICATIONS

- Capacity: ≥9995 endotoxin units (EU) removed by 1ml resin from 5ml test containing 10,000EU. ≥99.95% removal.
- Fractionation Range: 10-4,000kDa for proteins
- Bead Structure: 6% cross-linked agarose

## IMPORTANT INFORMATION

- Non specific binding of hydrophobic molecules may occur. To limit non-specific binding buffer all solutions to physiological pH. To inhibit ionic interactions, use a final concentration of 0.1-0.5M NaCl.
- Avoid the use of chaotropes and detergents as this interfere with binding to polymyxin sulfate.
- Some proteins, including BSA, bind tightly to endotoxins reducing their interaction and removal by the polymyxin sulfates. Increasing the volume of EndotoxinOUT™ resin to endotoxin containing sample may help.
- In some cases, the interaction between endotoxin and polymyxin sulfate is not inhibited, resulting in immobilization of the protein to the resin.
- Optimal performance of the EndotoxinOUT™ is best achieved under gravity flow chromatography, with an incubation period, allowing for more time for endotoxins to be in contact with polymyxin sulfate.

## ADDITIONAL ITEMS REQUIRED


- Columns (optional): G-Biosciences offers columns for a large range of resin volumes (Cat. # 786-718 to 786-724)
- Regeneration Solution: 1% sodium deoxycholate in endotoxin free water. Do not substitute a different detergent.
- Endotoxin-Free Water (Cat. # 786-670, 786-671)

## PROTOCOL

- Regenerate the EndotoxinOUT™ before and after every use.
  - Use endotoxin free solutions to prevent further endotoxin contamination.
  - Degas all solutions to prevent introducing air bubbles that inhibit column flow.
  - For 786-369, EndotoxinOUT™ columns, skip the degassing step.
1. Allow the resin and reagents to equilibrate to room temperature.
  2. Degas the EndotoxinOUT™ slurry by placing an appropriate volume in a suction filter flask with a magnetic stirrer. Apply a vacuum and gently stir the resin for 15 minutes.
  3. Add an appropriate volume of degassed, homogenous EndotoxinOUT™ resin to a suitable column and allow to settle for 30 minutes.
  4. *Regeneration:* Wash the resin with 5 resin volumes of Regeneration Solution, then wash with 5 resin volumes of Endotoxin-Free Water.
  5. Equilibrate the column with 5 resin volumes of an appropriate endotoxin-free buffer or water.
  6. Apply the sample to the column and add the bottom then top cap. Incubate at room temperature for 30-60 minutes.
  7. Elute the sample with appropriate aliquots of endotoxin-free buffer or water. Repeat the elution 3-6 times and monitor elutions by absorbance at 280nm or an appropriate assay.  
**NOTE:** Use extreme caution when handling samples to prevent contamination from poor handling, dirty glassware, etc.
  8. Repeat step 4 to regenerate the column and store in 25% ethanol at 4°C. The resin can be regenerated at least 10 times.

## RELATED PRODUCTS

Download our Sample Preparation Handbook.



<http://info.gbiosciences.com/complete-protein-sample-preparation-handbook/>

For other related products, visit our website at [www.GBiosciences.com](http://www.GBiosciences.com) or contact us.

Last saved: 6/12/2015 TN



[www.GBiosciences.com](http://www.GBiosciences.com)