

Rabbit IgG TrueBlot® Western Blot Kit Protocol - *Research Use Only*

Catalog No. [88-8886-31](#)

Materials Included

- Rabbit IgG TrueBlot: 50 µL. An HRP-conjugated second step reagent reacting with rabbit IgGs for optimal signal detection in immunoprecipitation/immunoblotting experiments (1000X) - [18-8816-31](#)
- TrueBlot® Enhancer Solution: 25 ml - 88-888C-ES
- TrueBlot® Blocker: 10 g - 88-888C-BC
- TrueBlot® Assay Buffer: 30 ml. 20X - 88-888C-AB
- TrueBlot® Substrate A: 12.5 ml - 88-888C-SA
- TrueBlot® Substrate B: 12.5 ml - 88-888C-SB

Materials Not Included

- Protein A agarose beads or Rabbit Ig IP beads (Cat. No. [00-8800-25](#)) which is included in the kit (Cat. No. [88-1688-31](#))
- Immunoprecipitation antibody
- PVDF or nitrocellulose membrane (0.2 of 0.45 µm)
- Immunoblotting rabbit primary antibody
- Chemiluminescent Substrate
- X-ray film

Buffers (see recipe below)

- Lysis buffer (with protease inhibitors)
- Cold PBS
- SDS-PAGE sample buffer with reducing agent
- Protease inhibitors
- Blocking Buffer
- TBST-T

Instruments

- Centrifuge
- Rocking platform or rotator
- SDS PAGE and Immunoblotting equipment and reagents

Step I: Preparation of Cell Lysate

1. Harvest approximately 1×10^7 cells by using cell scraper and transfer to conical tube. If working with adherent cells you can skip this step and lyse directly on the plate (see Step 6)
Note: The total number of cells per ml and the cell equivalent loaded per lane of gel should be optimized specifically for each protein and antibody. Alternatively, protein concentration can be determined using Bradford/Lowry or other protein assay.
2. Wash cells with ~10 mL of cold PBS and centrifuge at 400 xg for 10 minutes at 4°C.
3. Discard the supernatant and repeat step 2.
4. After the second wash, remove the supernatant and resuspend the cell pellet in 1 mL of cold Lysis Buffer containing protease Inhibitors (such as a cocktail- see recipe below). Final concentration of cells should be about 1×10^7 cells/mL.
Note: If using adherent cells, the cold Lysis Buffer can be added directly to the plate and put on a rocker at 4°C. Harvest by either scraping or collecting just the supernatant and proceed to Step 8.



5. Gently vortex/mix and transfer to 1.5 mL tube.
6. Place on ice for 30 minutes, with occasional mixing.
7. Centrifuge at 10,000 xg for 15 minutes at 4°C.
8. Carefully collect the supernatant, without disturbing the pellet and transfer to a new clean tube and discard pellet.
9. The protein concentration can be determined by Bradford or other assay. Samples can be diluted to ~1 g/.
10. The cell lysate can be frozen at this point for long-term storage at <-70°C.

Step II: Cell Lysate Preclearing

1. Resuspend the immobilized Protein A or Anti-Rabbit IgG bead slurry by gently vortexing. Remove 50 μ L and wash in Lysis buffer or IP buffer, if different. Resuspend in 50 μ L IP buffer.
2. Add 500 μ L of cell lysate (~ 5×10^6 cells or ~500 g/) to equilibrated beads slurry to and incubate on a rocking platform or a rotator for 30-60 minutes at 4°C.
3. Centrifuge at 2,500 xg for 2-3 minutes at 4°C and transfer the supernatant to a new 1.5 mL tube. If any of the bead slurry has been transferred, centrifuge again and carefully transfer the supernatant to another fresh 1.5 mL tube.

Step III: Immunoprecipitation

1. Add 1-10 μ g of immunoprecipitation antibody to the tube containing the cold precleared cell lysate.
Note: This concentration of monoclonal antibody is suggested as a starting point. Each investigator may desire to titrate the concentration of antibody and volume of cell lysate in preliminary experiments to determine the optimal conditions. e.g., 1-10 μ g/ 10^7 cells/1 mL lysate. Typically, 2 μ g is a sufficient amount of antibody to maximally immunoprecipitate most antigens in 1 ml of extract from 1×10^7 cells. Using as little IP antibody as possible minimizes potential contamination of SDS reduced sample with nonreduced immunoprecipitating antibody light chain. It is not recommended to use more than 10 μ g (per mL) or a final of 5 μ g per lane.
2. Incubate at 4°C for 1 hour on a rocking platform or a rotator.
3. Add at least 50 μ L of pre-equilibrated bead slurry to capture the immune complexes.
4. Incubate for 1 hour or overnight at 4°C on a rocking platform or a rotator.
Note: Step 1 and 3 can combined for a single incubation.
5. Centrifuge the tube at 2,500 xg for 30 seconds at 4°C.
6. Carefully remove supernatant completely and wash the beads 3-5 times with 500 μ L of cold Lysis Buffer, centrifuging to pellet beads in between each wash. In order to minimize background, care should be given to remove the supernatant completely after each wash.
7. After the last wash, carefully aspirate supernatant and add 50 μ L of 1X Laemmli sample buffer (or any equivalent SDS-PAGE sample loading Buffer) to bead pellet.
Note: Please take into consideration composition of the Loading buffer. Reducing agents can be added.
8. Vortex and heat to 90-100°C for 10 minutes.
9. Centrifuge at 10,000 xg for 5 minutes, collect supernatant carefully and load onto the gel.
10. Alternatively, the supernatant samples can be collected, transferred to clean tube and frozen at <-70°C if the gel is to be run later.
11. Follow manufacturer's instructions for SDS-PAGE.

Step IV: Immunoblotting (Western Blotting, WB)

1. Transfer proteins from the gel onto PVDF or nitrocellulose membrane following instructions provided by the transfer system manufacturer for best protein transfer results.
2. Optional: To determine whether the proteins have been transferred to membrane, stain with a 0.1% Ponceau S solution. Protein bands can be visualized after staining for 5 minutes. To remove the Ponceau S stain, rinse with distilled water or TBS-T until most of the dye is removed before moving on to blocking step. Residual dye will not affect subsequent steps
3. Remove membrane and soak in transfer buffer.
4. Under chemical hood, place the membrane in TrueBlot Enhancer Solution and soak for 2 minutes, then wash with TBS-T.



5. Place the membrane into the 5% TrueBlot Blocker in TrueBlot Assay Buffer (enough to cover the membrane) and incubate for 2 hours at room temperature or overnight at 4°C on a rocking platform. [Preparation of 5% TrueBlot Blocker in TrueBlot Assay Buffer: Dilute 20X TrueBlot Assay Buffer with dH₂O to 1X. Using TrueBlot Blocker Powder, make a 5% (w/v) solution.]
Note: it is recommended to use Milk as the blocking reagent as BSA does not effectively block the reduced Ig chain recognition.
6. Remove the blocking buffer and rinse blot with TBS-T.
7. Prepare the primary rabbit immunoblotting antibody in TrueBlot Blocker as recommended by the supplier. If the recommended concentration is not known use a standard concentration of 1-2 µg/ml. If using hybridoma tissue culture supernatant or serum for immunoblotting, preliminary experiments should be performed to evaluate whether dilution of the supernatant or serum is needed for best results.
8. Incubate the blot with primary antibody for at least 2 hours at room temperature or overnight at 4°C on rocking platform.
Note: Shorter times should be determined empirically for optimal results.
9. After the overnight incubation of the membrane with the primary antibody, wash the blot at least 3-5 times in TBS-T, each wash for a minimum of 5-10 minutes each. Total should be more than 1 hour.
10. Prepare the secondary antibody Rabbit IgG TrueBlot® at a 1:1,000 dilution in the Blocking Buffer.
Note: Please avoid the presence of sodium azide in this step as it is deleterious to the HRP enzyme.
11. Incubate the blot with the Trueblot secondary antibody for one hour at room temperature on a rocking platform.
12. Wash the blot at least 3-5 times in TBS-T, each wash for at least 5 minutes each.
13. Prepare Substrate: Mix equal volumes of Substrate A and B
14. Incubate the blot in chemiluminescent-HRP substrate working solution (combined A and B) for 1-5 minutes.
15. Expose the blot to X-ray film for an appropriate time period. For best results, expose for ten seconds, one minute, five minutes and 20 minutes.

Related Products

Chemiluminescent FemtoMax™ Super Sensitive HRP Substrate for Microwell/Membrane (2 component system) - [FEMTOMAX-020](#)

BLOTTO Immunoanalytical Grade (Non-Fat Dry Milk) - [B501-0500](#)

10X TTBS pH 7.5 (1.0 M Tris HCl 1.5 M Sodium Chloride 0.1% (w/v) Tween-20) - [MB-013](#)

2X SDS-PAGE Sample Buffer without DTT or β-MeOH (125 mM Tris HCl, 20% Glycerol, 4% SDS, 0.1% bromphenol blue, pH 6.8) - [MB-018](#)

Related Phospho Specific Primary Antibodies

TrueBlot® is ideal for studying post-translational protein modifications, including phosphorylation, or protein-protein interactions. Our portfolio offers a large and diverse set of [phospho-specific antibodies](#) and reagents that detect phosphorylated targets with great precision. Additionally, Rockland [phospho antibodies](#) provide accuracy and dependability that clients within the life sciences have come to expect.

- Anti-AKT pS473 (Mouse) [200-301-268](#)
- Anti-AKT pT308 (Mouse) [200-301-269](#)
- Anti-ATM Protein Kinase pS1981 (Mouse) [200-301-400](#)
- Anti-Histone H2AvD pS137 (Rabbit) [600-401-914](#)
- Anti-Myosin pS19/pS20 (Rabbit) [600-401-416](#)
- Anti-SMAD3 pS423 pS425 (Rabbit) [600-401-919](#)



Solutions & Recipes

2X SDS Reducing Sample Loading Buffer (containing 50 mM DTT)

- 950 mL of 2X SDS sample buffer
- 50 mL of 1M DTT

Note: Use within 1 hour and discard remainder.

2X SDS Sample Buffer

- 6% SDS
- 25mM Tris base pHed to 6.5 with HCl
- 10% glycerol
- Bromphenol blue

Note: Can be stored long term at -20°C and for up to 1 month at room temperature.

1M DTT

- Can be made fresh or can be stored as aliquots at -20°C for 6 months or at 4°C for 2 weeks. Avoid repeated freeze thaws.

TBS-Tween (TBS-T)

- 25 mM Tris-HCl, pH 8.0
- 125 mM NaCl
- 0.1% Tween 20

Blocking Buffer:

- 5% Carnation nonfat dry milk in TBS-T

Note: Milk solution should be stored at 4°C short term or -20°C for long term.

NP-40 Cell Lysis Buffer:

- 50mM Tris-HCl pH 8.0
- 150mM NaCl
- 1% NP-40

RIPA Buffer:

- 50mM Tris-HCl pH 7.4
- 1% NP-40
- 0.25% Na-deoxycholate
- 150mM NaCl
- 1mM EDTA

Protease Inhibitor Cocktail (100X):

- PMSF, 5mg (50µg/ml)
- Aprotinin, 100µg (1µg/ml)
- Leupeptin, 100µg (1µg/ml)
- Pepstatin, 100µg (1µg/ml)

Phosphatase Inhibitor (100X):

- 1mM Na₃VO₄
- 1mM NaF



Rabbit TrueBlot® Troubleshooting Chart

Problem	Possible Cause	Solution
A. No Signal	<ol style="list-style-type: none"> 1. Weak primary antibody 2. NaN₃ is present during HRP-substrate incubation 3. Primary antibody is not a rabbit IgG 4. Target protein is not expressed in the sample or present at very low level 5. Antigen is present in blocking solution 	<ol style="list-style-type: none"> 1. Use only primary antibodies optimized for immunoblotting 2. Incubate HRP-substrate in NaN₃ free buffer 3. Use only rabbit IgG as primary antibody for Rabbit IgG TrueBlot® 4. Use as positive control, sample known to contain the target protein and optimize the amount of protein loaded 5. Change blocking reagents
B. High background	<ol style="list-style-type: none"> 1. Non-optimized primary antibody 2. Insufficient washing 3. Membrane was allowed to dry and not re-wetted 4. Insufficient blocking 	<ol style="list-style-type: none"> 1. Use only primary antibodies optimized for immunoblotting 2. Increase volume, number and duration of washes; increase salt content of the wash buffer (see Appendix) 3. Ensure membrane is not dried during immunoblotting procedures. Immobilon-P and other PVDF membranes must be wetted in methanol and equilibrated in buffer 4. 5% (w/v) nonfat dry milk is the best blocking agent. BSA is specifically not recommended.
C. I see Ig in addition to my specific band of interest	<ol style="list-style-type: none"> 1. Improper sample preparation 	<ol style="list-style-type: none"> 1. Follow sample preparation procedure
D. I see other bands in addition to my specific band of interest	<ol style="list-style-type: none"> 1. Poor primary antibody: low signal/high noise 	<ol style="list-style-type: none"> 1. Use primary antibodies optimized for immunoblotting (high signal/noise) 2. Possible different isoforms/modifications of the protein of interest

